

Renal Leiomyoma in a 54-Year-Old Woman

**AUTHORS:**

ME, C, P, I, A

**CORRESPONDING AUTHOR:**

Mary E. Tabatneck, MD

35

A 40

Boston, MA 0244

ETab@bwh.harvard.edu

**AUTHOR AFFILIATION:**

Dartmouth-Hitchcock Medical Center

122Q

Lebanon, NH 03756

<b>Background</b>	We present a case of a clinically symptomatic 54-year-old woman with a 16 cm renal leiomyoma diagnosed with histopathology after radical nephrectomy.
<b>Summary</b>	Our patient is a 54-year-old woman who presented with a palpable left abdominal mass. The patient underwent radical nephrectomy with histopathologic evaluation revealing renal leiomyoma. Renal leiomyomas are extremely rare neoplasms most commonly diagnosed by histopathologic evaluation. Imaging of large renal masses fails to establish diagnosis since large benign masses such as leiomyomas may present with necrosis, hemorrhage, or other symptoms concerning of malignancy.
<b>Conclusion</b>	While most renal leiomyomas are incidentally identified on imaging, renal leiomyomas rarely can present as a large symptomatic mass. Often symptomatic masses demonstrate malignant features on imaging, highlighting the importance of histopathologic diagnosis.
<b>Keywords</b>	renal leiomyoma; nephrectomy

**DISCLOSURE STATEMENT:**

The authors indicated no potential conflicts of interest.


**FUNDING/SUPPORT:**

The authors indicated no potential conflicts of interest.

RECEIVED: December 15, 2020

REVISION RECEIVED: February 5, 2021

ACCEPTED FOR PUBLICATION: March 25, 2021




## Case Discussion


Renal leiomyomas are benign neoplasms of mesenchymal origin, typically presenting as incidental discoveries on autopsy. Having an incidence of only 5% in autopsy specimens, renal leiomyomas are rare entities, and their clinical incidence is thought to be much lower.<sup>1-3</sup> Renal leiomyomas lack distinguishing clinical or radiographic characteristics, and diagnosis is typically made after surgical excision.<sup>4,5</sup> We report a case of a large renal leiomyoma in a 54-year-old woman that was surgically excised with radical nephrectomy.

The patient is a 54-year-old woman who presented to the clinic with a one-year history of left-sided abdominal discomfort, fullness, and distension. She noted occasional constipation over the year but denied fevers, night sweats, unexpected weight loss, worsening fatigue, nausea, or vomiting. A palpable mass was appreciated on physical exam. Laboratory values on her complete blood count and comprehensive metabolic panel were unremarkable. Her primary care physician obtained an ultrasound demonstrating a heterogenous retroperitoneal mass measuring 15 × 9 × 14 cm arising from the left adrenal gland or left kidney. Subsequent magnetic resonance imaging of the abdomen performed at an outside institution (Figure 1) confirmed a 12 × 9 × 16 cm left-sided retroperitoneal mass displacing the left kidney and the pancreas. Although the mass was well-circumscribed, the origin of the mass was unclear. Intralésional regions of low intensity suggested central necrosis.

The patient was referred to the surgical oncology service for preoperative evaluation and operative planning for the excision of the mass. Given the size and symptomatic nature of the mass, a decision was made to pursue a more radical surgical excision instead of a percutaneous biopsy. Preoperative computed tomography of the abdomen and pelvis was obtained to assess the anatomic planes for resection because the MRI images performed at the previous medical facility were not readily available (Figure 2).



Axial at its largest dimensions; images were used for preoperative planning and visualization of planes of dissection.



Coronal at its largest dimensions; images were used for preoperative planning and visualization of planes of dissection.

The patient underwent an exploratory laparotomy and was found to have a large left renal tumor. Intraoperative ultrasound was used to delineate the borders of the mass. No obvious pancreatic connections were appreciated on ultrasound. A radical left nephrectomy was performed, and one lymph node that appeared as a mass near the ligament of Treitz was removed. She tolerated the procedure well without intraoperative complications. Her postoperative course was uncomplicated. Pathology of the resected mass confirmed a well-circumscribed leiomyoma without atypia, necrosis, or mitotic activity, originating from the renal capsule. The tumor was positive for actin, desmin, smooth muscle myosin, estrogen receptor, and progesterone receptor. The tumor was negative for CD34, c-Kit, and DOG-1. Resected lymph node sent with the specimen was benign. She was seen in clinic at her four-week follow-up and was doing well without late postoperative complications or evidence of worsening kidney function.

#### D. c, f, g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z

Leiomyomas may arise from smooth muscle cells anywhere along the genitourinary tract. Renal leiomyomas are a subset that most commonly arises from the subcapsular tissue but may also arise from the capsule or renal pelvis.<sup>1,4</sup> Although renal leiomyomas are identified in approximately 4.2%-5.2% of autopsy specimens, the clinical incidence of renal leiomyomas is currently unknown.<sup>5</sup> In a review of 1030 nephrectomies performed over a ten-year period, renal leiomyomas accounted for 0.29% of treated renal masses and 1.5% of benign renal masses.<sup>6</sup> A 2015 study evaluating approximately 4000 renal neoplasms over a 22-year period identified four renal leiomyomas, suggesting an incidence of 0.001%.<sup>4</sup> A case series involving three patients and associated literature review suggested that symptomatic renal leiomyomas may have a predilection for females in their second to fifth decades of life.<sup>6</sup> The average size of asymptomatic renal leiomyomas is approximately 5 mm, while the average size of symptomatic renal leiomyomas is approximately 12.3 cm.<sup>7,8</sup>

The differential diagnosis for renal tumors includes benign masses such as renal leiomyomas and angiomyolipomas as well as malignant tumors such as leiomyosarcomas and renal cell carcinomas. Renal leiomyomas have no specific clinical finding and thus would share common clinical characteristics with any of these masses. Large retroperitoneal tumors generally present with palpable flank mass or flank pain along with mass effects on local structures.<sup>2,3</sup> These symptoms are non-specific. Hematuria has been

